Guideline for Stroke Prevention in Patients with Atrial Fibrillation (AF)

Definition

An irregularly irregular pulse, confirmed by an ECG that demonstrates an absence of P waves

Background

- May be chronic, paroxysmal, or persistent
- For the purposes of thromboembolic prophylaxis, this guideline also applies to the management of atrial flutter
- Prevalence increases steadily with age: 0.5% age 50-59; 8.8% age 80-89
- Commonest arrhythmia requiring hospitalisation
- AF is associated with a fivefold increase in risk of ischaemic stroke
- CT stùdies have shown that silent cerebral infarction is present in 26% of patients with AF

Presentation/symptoms

- Palpitations
- Shortness of breath
- Chest pain
- Lethargy
- Systemic embolisation
- Opportunistic taking of pulse

Causes of AF

- Ischaemic heart disease
- Hypertension
- Valvular heart disease
- Thyrotoxicosis
- Infection

DENTIFY

Idiopathic

Box 1: When to refer for Cardiology Opinion (

- AF with broad complexes on ECG
- AF resistant to 'usual' drugs for rate control
- Recent onset
- Heart failure
- Frequent attacks of paroxysmal AF
- Syncope related to paroxysmal AF
- Heart murmur

UNLESS EXCLUDED BY
BOX 1, ASSESS
THROMBOEMBOLIC RISK
AND DETERMINE
MANAGEMENT PLAN
USING CHAD2 SCORE
(BELOW)

CHAD2 Score (A)


22	С	Congestive heart failure
ASSESS	Н	History of hypertension
ó	A	Age > 75 years
1	D	Diabetes
	2	Prior stroke or TIA

Total Score

CHAD2 Score	Adjusted Annual Stroke Rate (%)	NNTs	Risk of major bleed (per 100 patient years)	
0 1	1.9 2.8	80 55	Aspirin	1.5
2 3 4 5 6	4.0 5.9 8.5 12.5 18.2	38 26 18 12 8	Warfarin	2.2

PLEASE NOTE:
BENEFIT OF
WARFARIN
OUTWEIGHS RISK
WHEN CHAD2
SCORE ≥ 2

≥ = 'greater than or equal to'



- Medical eg. bleeding lesions, liver disease, uncontrolled hypertension
- Mental or physical frailty i.e. falls, confusion
- Non-compliance i.e. with drug or clinic
- Patient refusal

Read codes	4 byte	5.byte	Read 3
Atrial fibrillation	G67.	G5730	G5730
Warfarin	Bs1.	Bs1	x01o4
ECG: general	321.	321	X77As
Aspirin prophylaxis - IHD	8B63	8B63.	XM1Xx
Aspirin contraindicated		8124	XaDvH
INR	42QE	42QE.	42QE.

References

CHAD2: JAMA. 2001; 285 (22): 2864-2870 Risk of bleeding: NEJM. 2001; 20 (345):

1444-1451

Clopidogrel: Lancet. 1996; 348: 1329-1339

INRs: BMJ. 1997; 317: 1563

Inclusion of atrial flutter and paroxysmal AF:

EBM. 2001; 6: 150

Grade of evidence: (A = systematic review or RCT; B = robust experimental or observational study; 6 = expert committee or local opinion